

EXTREME VALUES OF $|\zeta(1 + it)|$

ANDREW GRANVILLE AND K. SOUNDARARAJAN

To Professor K. Ramachandra on the occasion of his seventieth birthday

1. INTRODUCTION

Improving on a result of J.E. Littlewood, N. Levinson [3] showed that there are arbitrarily large t for which $|\zeta(1 + it)| \geq e^\gamma \log_2 t + O(1)$. (Throughout $\zeta(s)$ is the Riemann-zeta function, and $\log_j$ denotes the j -th iterated logarithm, so that $\log_1 n = \log n$ and $\log_j n = \log(\log_{j-1} n)$ for each $j \geq 2$.) The best upper bound known is Vinogradov's $|\zeta(1 + it)| \ll (\log t)^{2/3}$.

Littlewood had shown that $|\zeta(1 + it)| \lesssim 2e^\gamma \log_2 t$ assuming the Riemann Hypothesis, in fact by showing that the value of $|\zeta(1 + it)|$ could be closely approximated by its Euler product for primes up to $\log^2(2 + |t|)$ under this assumption. Under the further hypothesis that the Euler product up to $\log(2 + |t|)$ still serves as a good approximation, Littlewood conjectured that $\max_{|t| \leq T} |\zeta(1 + it)| \sim e^\gamma \log_2 T$, though later he wrote in [5] (in connection with a q -analogue): “*there is perhaps no good reason for believing ... this hypothesis*”.

Our Theorem 1 evaluates the frequency with which such extreme values are attained; and if this density function were to persist to the end of the viable range then this implies the conjecture that

$$(1.1a) \quad \max_{t \in [T, 2T]} |\zeta(1 + it)| = e^\gamma (\log_2 T + \log_3 T + C_1 + o(1)),$$

for some constant C_1 . In fact it may be that $C_1 = C + 1 - \log 2$, where

$$C = \int_0^2 \log I_0(t) \frac{dt}{t^2} + \int_2^\infty (\log I_0(t) - t) \frac{dt}{t^2},$$

and $I_0(t) := \mathbb{E}(e^{\operatorname{Re}(tX)}) = \sum_{n=0}^\infty (t/2)^{2n}/n!$ is the Bessel function (with X a random variable equidistributed on the unit circle). In Theorem 2 we show that there are arbitrarily large t for which $|\zeta(1 + it)| \geq e^\gamma (\log_2 t + \log_3 t - \log_4 t + O(1))$, which improves upon Levinson's result but falls a little short of our conjecture.

Le premier auteur est partiellement soutenu par une bourse de la Conseil de recherches en sciences naturelles et en génie du Canada. The second author is partially supported by the National Science Foundation.

Levinson also showed that $1/|\zeta(1+it)| \geq \frac{6e^\gamma}{\pi^2}(\log_2 t - \log_3 t + O(1))$ for arbitrarily large t . Theorem 1 exhibits even smaller values of $|\zeta(1+it)|$ and determines their frequency. Extrapolating Theorem 1 we are also led to conjecture that

$$(1.1b) \quad \max_{t \in [T, 2T]} 1/|\zeta(1+it)| = \frac{6e^\gamma}{\pi^2}(\log_2 T + \log_3 T + C_1 + o(1));$$

but only succeed in proving that $1/|\zeta(1+it)| \geq \frac{6e^\gamma}{\pi^2}(\log_2 t - O(1))$ for arbitrarily large t . K. Ramachandra [6] has obtained results analogous to Levinson's in short intervals, and R. Balasubramanian, Ramachandra and A. Sankaranarayanan [1] have considered extreme values of $|\zeta(1+it)|^{e^{i\theta}}$ for any $\theta \in [0, 2\pi)$.

To be more precise let us define, for $T, \tau \geq 1$,

$$\begin{aligned} \Phi_T(\tau) &:= \frac{1}{T} \text{meas}\{t \in [T, 2T] : |\zeta(1+it)| > e^\gamma \tau\}, \\ \text{and } \Psi_T(\tau) &:= \frac{1}{T} \text{meas}\{t \in [T, 2T] : |\zeta(1+it)| < \frac{\pi^2}{6e^\gamma \tau}\}. \end{aligned}$$

Theorem 1. *Let T be large. Uniformly in the range $1 \ll \tau \leq \log_2 T - 20$ we have*

$$\Phi_T(\tau) = \exp\left(-\frac{2e^{\tau-C-1}}{\tau}\left(1 + O\left(\frac{1}{\tau^{\frac{1}{2}}} + \left(\frac{e^\tau}{\log T}\right)^{\frac{1}{2}}\right)\right)\right),$$

where c is a positive constant. The same asymptotic also holds for $\Psi_T(\tau)$.

With a judicious application of the pigeonhole principle we can exhibit even larger values of $|\zeta(1+it)|$, indeed of almost the same quality as the conjectured (1.1a).

Theorem 2. *For large T the subset of points $t \in [0, T]$ such that*

$$|\zeta(1+it)| \geq e^\gamma(\log_2 T + \log_3 T - \log_4 T - \log A + O(1))$$

has measure at least $T^{1-\frac{1}{A}}$, uniformly for any $A \geq 10$.

One can also establish results analogous to Theorems 1 and 2 for the distribution of values of $|L(1, \chi)|$ where χ ranges over all non-trivial characters modulo a large prime p (see section 7 for further details). In fact Theorems 1 and 2 hold almost verbatim, just changing T to p . If one also averages over p in a dyadic interval $P \leq p \leq 2P$ then one can obtain asymptotics for the distribution function in the wider range $1 \ll \tau \leq \log_2 P + \log_3 P - O(1)$ (which we expect is the full range, up to the explicit value of the “ $O(1)$ ”).

As in [2] we can compare the distribution of $\zeta(1+it)$ with that of an appropriate probabilistic model. Let $X(p)$ denote independent random variables uniformly distributed on the unit circle, for each prime p . We extend X multiplicatively to all integers n : that is set $X(n) = \prod_{p^\alpha \parallel n} X(p)^\alpha$. We wish to compare the distribution of values of $\zeta(1+it)$ with the distribution of values of the random Euler products $L(1, X) := \prod_p (1 - X(p)/p)^{-1}$ (these products converge with probability 1). Now define

$$\Phi(\tau) = \text{Prob}(|L(1, X)| \geq e^\gamma \tau) \quad \text{and} \quad \Psi(\tau) = \text{Prob}(|L(1, X)| \leq \frac{\pi^2}{6e^\gamma \tau}).$$

By the same methods one can show that $\Phi(\tau)$ and $\Psi(\tau)$ satisfy the same asymptotic as $\Phi_T(\tau)$ as in Theorem 1, but for arbitrary τ (see the remarks immediately after the proof of Theorem 1).

2. PRELIMINARIES

We collect here some standard facts on $\zeta(s)$ which will be used later.

Lemma 2.1. *Let $y \geq 2$ and $|t| \geq y+3$ be real numbers. Let $\frac{1}{2} \leq \sigma_0 < 1$ and suppose that the rectangle $\{z : \sigma_0 < \operatorname{Re}(z) \leq 1, | \operatorname{Im}(z) - t| \leq y+2\}$ is free of zeros of $\zeta(z)$. Then for any $\sigma_0 < \sigma \leq 2$ and $|\xi - t| \leq y$ we have*

$$|\log \zeta(\sigma + i\xi)| \ll \log |t| \log(e/(\sigma - \sigma_0)).$$

Further for $\sigma_0 < \sigma \leq 1$ we have

$$\log \zeta(\sigma + it) = \sum_{n=2}^y \frac{\Lambda(n)}{n^{\sigma+it} \log n} + O\left(\frac{\log |t|}{(\sigma_1 - \sigma_0)^2} y^{\sigma_1 - \sigma}\right),$$

where we put $\sigma_1 = \min(\sigma_0 + \frac{1}{\log y}, \frac{\sigma + \sigma_0}{2})$.

Proof. The first assertion follows from Theorem 9.6(B) of Titchmarsh [8]. In proving the second assertion we may plainly suppose that $y \in \mathbb{Z} + \frac{1}{2}$. Then Perron's formula gives, with $c = 1 - \sigma + \frac{1}{\log y}$,

$$\begin{aligned} \frac{1}{2\pi i} \int_{c-iy}^{c+iy} \log \zeta(\sigma + it + w) \frac{y^w}{w} dw &= \sum_{n=2}^y \frac{\Lambda(n)}{n^{\sigma+it} \log n} + O\left(\frac{1}{y} \sum_{n=1}^{\infty} \frac{y^c}{n^{\sigma+c}} \frac{1}{|\log(y/n)|}\right) \\ (2.1) \qquad \qquad \qquad &= \sum_{n=2}^y \frac{\Lambda(n)}{n^{\sigma+it} \log n} + O(y^{-\sigma} \log y). \end{aligned}$$

We now move the line of integration to the line $\operatorname{Re}(w) = \sigma_1 - \sigma < 0$. Our hypothesis ensures that the integrand is regular over the region where the line is moved, except for a simple pole at $w = 0$ which leaves the residue $\log \zeta(\sigma + it)$. Thus the left side of (2.1) equals $\log \zeta(\sigma + it)$ plus

$$\frac{1}{2\pi i} \left(\int_{c-iy}^{\sigma_1 - \sigma - iy} + \int_{\sigma_1 - \sigma - iy}^{\sigma_1 - \sigma + iy} + \int_{\sigma_1 - \sigma + iy}^{c+iy} \right) \log \zeta(\sigma + it + w) \frac{y^w}{w} dw \ll \frac{\log |t|}{(\sigma_1 - \sigma_0)^2} y^{\sigma_1 - \sigma},$$

upon using the first part of the Lemma.

Using Lemma 2.1 we shall show that most of the time we may approximate $\zeta(s)$ by a short Euler product.

Lemma 2.2. *Let $\frac{1}{2} < \sigma \leq 1$ be fixed and let T be large. Let $T/2 \geq y \geq 3$ be a real number. The asymptotic*

$$\log \zeta(\sigma + it) = \sum_{n=2}^y \frac{\Lambda(n)}{n^{\sigma+it} \log n} + O(y^{(\frac{1}{2}-\sigma)/2} \log^3 T)$$

holds for all $t \in (T, 2T)$ except for a set of measure $\ll T^{5/4-\sigma/2} y (\log T)^5$.

Proof. This follows upon using the zero-density result $N(\sigma_0, T) \ll T^{3/2-\sigma_0} (\log T)^5$ (see Theorem 9.19 A of [8]) and appealing to Lemma 2.1 (taking $\sigma_0 = (1/2 + \sigma)/2$ there).

3. APPROXIMATING $\zeta(1+it)$ BY A SHORT EULER PRODUCT

Lemma 3.1. *Suppose $2 \leq y \leq z$ are real numbers. Then for arbitrary complex numbers $x(p)$ we have*

$$\frac{1}{T} \int_T^{2T} \left| \sum_{y \leq p \leq z} \frac{x(p)}{p^{it}} \right|^{2k} \ll \left(k \sum_{y \leq p \leq z} |x(p)|^2 \right)^k + T^{-\frac{2}{3}} \left(\sum_{y \leq p \leq z} |x(p)| \right)^{2k}$$

for all integers $1 \leq k \leq \log T / (3 \log z)$.

Proof. The quantity we seek to estimate is

$$\sum_{\substack{p_1, \dots, p_k \\ y \leq p_j \leq z}} \sum_{\substack{q_1, \dots, q_k \\ y \leq q_j \leq z}} \overline{x(p_1) \cdots x(p_k)} x(q_1) \cdots x(q_k) \frac{1}{T} \int_T^{2T} \left(\frac{p_1 \cdots p_k}{q_1 \cdots q_k} \right)^{it} dt.$$

The diagonal terms $p_1 \cdots p_k = q_1 \cdots q_k$ contribute

$$\ll k! \left(\sum_{y \leq p \leq z} |x(p)|^2 \right)^k.$$

If $p_1 \cdots p_k \neq q_1 \cdots q_k$ then as both quantities are below $z^k \leq T^{\frac{1}{3}}$ we have that

$$\frac{1}{T} \int_T^{2T} \left(\frac{p_1 \cdots p_k}{q_1 \cdots q_k} \right)^{it} dt \ll \frac{1}{T |\log(p_1 \cdots p_k / q_1 \cdots q_k)|} \ll T^{-\frac{2}{3}}.$$

Hence the off diagonal terms contribute $\ll T^{-\frac{2}{3}} (\sum_{y \leq p \leq z} |x(p)|)^{2k}$, proving the Lemma.

Define $\zeta(s; y) := \prod_{p \leq y} (1 - p^{-s})^{-1}$.

Proposition 3.2. *Let T be large and let $\log T (\log_2 T)^4 \geq y \geq e^2 \log T$ be a real number. Then there is a positive constant c such that*

$$\zeta(1+it) = \zeta(1+it; y) \left(1 + O\left(\frac{\sqrt{\log T}}{\sqrt{y} \log_2 T} \right) \right)$$

for all $t \in (T, 2T)$ except for a set of measure at most $T \exp(-\log T / 50 \log_2 T)$.

Proof. Setting $z = (\log T)^{100}$ we deduce from Lemma 2.2 that $\zeta(1+it) = \zeta(1+it; z) (1 + O(1/\log T))$ for all $t \in (T, 2T)$ except for a set of measure at most $T^{4/5}$. Using Lemma 3.1 with $k = \lfloor \log T / (300 \log_2 T) \rfloor$ and $x(p) = 1/p$ we get that

$$\begin{aligned} \frac{1}{T} \int_T^{2T} \left| \sum_{y \leq p \leq z} \frac{1}{p^{1+it}} \right|^{2k} dt &\ll \left(k \sum_{y \leq p \leq z} \frac{1}{p^2} \right)^k + T^{-\frac{2}{3}} \left(\sum_{y \leq p \leq z} \frac{1}{p} \right)^{2k} \\ &\ll \left(\frac{\log T}{y} \right)^k \left(\frac{1}{10 \log y} \right)^{2k} + T^{-\frac{1}{3}}, \end{aligned}$$

and so

$$\left| \sum_{y \leq p \leq z} \frac{1}{p^{1+it}} \right| \leq \frac{\sqrt{\log T}}{\sqrt{y} \log y}$$

for all $t \in [T, 2T]$ except for a set of measure $\leq T \exp(-\log T/49 \log_2 T)$. The Proposition thus follows, by combining the above estimates, since

$$\zeta(1+it; y) = \zeta(1+it; z) \exp \left(- \sum_{y \leq p \leq z} \left(\frac{1}{p^{1+it}} + O\left(\frac{1}{p^2}\right) \right) \right).$$

4. MOMENTS OF SHORT EULER PRODUCTS

In this section we show how to evaluate large moments of the short Euler products obtained in §3.

Theorem 4.1. *Let $\log T(\log_2 T)^4 \geq y \geq e^2 \log T$ be a real number. Let $z = \delta k$ where $\delta = \pm 1$ and $2 \leq k \leq \log T/(e^{10} \log(y/\log T))$ is an integer. Then*

$$\begin{aligned} \frac{1}{T} \int_T^{2T} |\zeta(1+it; y)|^{2z} &= \sum_{\substack{n=1 \\ p|n \implies p \leq y}}^{\infty} \frac{d_z(n)^2}{n^2} \left(1 + O\left(\exp\left(-\frac{\log T}{2(\log_2 T)^4} \right) \right) \right) \\ &= \prod_{p \leq k} \left(1 - \frac{\delta}{p} \right)^{-2k\delta} \exp \left(\frac{2k}{\log k} \left(C + O\left(\frac{k}{y} + \frac{1}{\log k} \right) \right) \right). \end{aligned}$$

Throughout this section let z, y, k, δ be as in Theorem 4.1. If $k \leq 10^6$ then we divide $[1, y]$ into the intervals $I_0 = [k, y]$ and $I_1 = [1, k]$ and take here $J := 1$. If $k > 10^6$ then we define $J := [4 \log_2 k / \log 2] + 1$ and divide $[1, y]$ into the $J+1$ -intervals $I_0 = [k, y]$, $I_j = [k/2^j, k/2^{j-1})$ for $1 \leq j \leq J-1$, and $I_J = [1, k/2^J] \subset [1, k/(\log k)^4]$. Given a subset R of the index set $\{0, 1, \dots, J\}$ we define $\mathcal{S}(R)$ to be the set of integers n whose prime factors all lie in $\cup_{r \in R} I_r$. We also define

$$\zeta(s; R) := \prod_{p \in \cup_{r \in R} I_r} \left(1 - \frac{1}{p^s} \right)^{-1} = \sum_{n \in \mathcal{S}(R)} \frac{1}{n^s}.$$

Proposition 4.2. *Let R be any subset of $\{0, \dots, J\}$. Then we have that*

$$\frac{1}{T} \int_T^{2T} |\zeta(1+it; R)|^{2z} dt = \sum_{n \in \mathcal{S}(R)} \frac{d_z(n)^2}{n^2} \left(1 + O\left(\exp\left(-\frac{\log T}{2(\log_2 T)^4} \right) \right) \right).$$

Note that the first part of Theorem 4.1 follows from the case $R = \{0, 1, \dots, J\}$. While this is the case of interest for us, the formulation of Proposition 4.2 is convenient for our proof which is based on induction on the cardinality of R .

Lemma 4.3. *For any prime p we have*

$$\sum_{a=0}^{\infty} \frac{d_z(p^a)^2}{p^{2a}} = I_0\left(\frac{2k}{p}\right) \exp(O(k/p^2)).$$

Also

$$\left(1 - \frac{\delta}{p}\right)^{-2k\delta} \geq \sum_{a=0}^{\infty} \frac{d_z(p^a)^2}{p^{2a}} \geq \frac{1}{50} \min\left(1, \frac{p}{k}\right) \left(1 - \frac{\delta}{p}\right)^{-2k\delta},$$

so that if $\mathcal{P}$ is any subset of the primes $\leq y$ then, uniformly,

$$\sum_{\substack{n \geq 1 \\ p|n \implies p \in \mathcal{P}}} \frac{d_z(n)^2}{n^2} \geq T^{O(1/\log_2 T)} \prod_{p \in \mathcal{P}} \left(1 - \frac{\delta}{p}\right)^{-2k\delta}.$$

Proof. Since

$$\sum_{a=0}^{\infty} \frac{d_z(p^a)^2}{p^{2a}} = \int_0^1 \left|1 - \frac{e(\theta)}{p}\right|^{-2z} d\theta = \int_0^1 \exp(O(k/p^2)) \exp\left(2\frac{z}{p} \cos(2\pi\theta)\right) d\theta$$

we obtain the first assertion. The upper bound in the second statement follows since $|1 - e(\theta)/p|^{-\delta} \leq (1 - \delta/p)^{-\delta}$. When $p > k$ we have that $(1 - \delta/p)^{-2k\delta} \leq (1 - 1/\max(2, k))^{-2k} \leq 16$ and so the lower bound follows in this case. When $p \leq k$ consider only θ such that $e(\theta)$ lies on the arc $(\delta e^{-ip/(10k)}, \delta e^{ip/(10k)})$. For such θ we may check that $|1 - e(\theta)/p|^{-2k\delta} \geq (1 - \delta/p)^{-2k\delta} (1 - 1/(25k))^k \geq \frac{4}{5} (1 - \delta/p)^{-2k\delta}$ from which the lower bound in this case follows.

Now

$$\prod_{\substack{k < p \leq y \\ p \in \mathcal{P}}} \left(1 - \frac{\delta}{p}\right)^{-2k\delta} \leq \exp\left(O\left(\sum_{k < p \leq y} \frac{k}{p}\right)\right) \ll \left(\frac{\log y}{\log k}\right)^{O(k)} \ll T^{O(1/\log_2 T)},$$

and

$$\sum_{\substack{n \geq 1 \\ p|n \implies p \in \mathcal{P}}} \frac{d_z(n)^2}{n^2} > \sum_{\substack{n \geq 1 \\ p|n \implies p \leq k \text{ and } p \in \mathcal{P}}} \frac{d_z(n)^2}{n^2} \geq \prod_{\substack{p \leq k \\ p \in \mathcal{P}}} \frac{p}{50k} \left(1 - \frac{\delta}{p}\right)^{-2k\delta},$$

which together imply the third assertion by the prime number theorem.

Lemma 4.4. *Suppose $0 \leq r \leq J$ and put $M_0 := T^{\frac{1}{5}}$ and $M_r = T^{\frac{1}{5r+2}}$ for $r \geq 1$. Then we have that*

$$\sum_{\substack{m \in \mathcal{S}(\{r\}) \\ m \geq M_r}} \frac{2^{\omega(m)}}{m} \sum_{\ell \in \mathcal{S}(\{r\})} \frac{|d_z(m\ell)d_z(\ell)|}{\ell^2} \leq \left(\sum_{\ell \in \mathcal{S}(\{r\})} \frac{d_z(\ell)^2}{\ell^2}\right) \exp\left(-\frac{\log T}{(\log_2 T)^4}\right).$$

Proof. Denote the left side of the estimate in Lemma 4.4 by N_r and let

$$D_r = \sum_{\ell \in \mathcal{S}(\{r\})} \frac{d_z(\ell)^2}{\ell^2}.$$

For any $1 \geq \alpha > 0$ we have

$$\begin{aligned} N_r &\leq M_r^{-\alpha} \sum_{m \in \mathcal{S}(\{r\})} \frac{2^{\omega(m)}}{m^{1-\alpha}} \sum_{\ell \in \mathcal{S}(\{r\})} \frac{|d_z(m\ell)d_z(\ell)|}{\ell^2} \\ (4.1) \quad &= M_r^{-\alpha} \prod_{p \in I_r} \left(\sum_{a=0}^{\infty} \frac{|d_z(p^a)|^2}{p^{2a}} + 2 \sum_{u=1}^{\infty} \frac{1}{p^{u(1-\alpha)}} \sum_{a=0}^{\infty} \frac{|d_z(p^a)d_z(p^{u+a})|}{p^{2a}} \right). \end{aligned}$$

We record two bounds for the p th term of the product in (4.1): Firstly

$$\begin{aligned} \sum_{a=0}^{\infty} \frac{|d_z(p^a)|^2}{p^{2a}} + 2 \sum_{u=1}^{\infty} \frac{1}{p^{u(1-\alpha)}} \sum_{a=0}^{\infty} \frac{|d_z(p^a)d_z(p^{u+a})|}{p^{2a}} &\leq 2 \sum_{a=0}^{\infty} \frac{|d_z(p^a)|}{p^{a(1+\alpha)}} \sum_{u=-a}^{\infty} \frac{|d_z(p^{u+a})|}{p^{(u+a)(1-\alpha)}} \\ (4.2) \quad &= 2 \left(1 - \frac{\delta}{p^{1-\alpha}}\right)^{-\delta k} \left(1 - \frac{\delta}{p^{1+\alpha}}\right)^{-\delta k}. \end{aligned}$$

Secondly, since $|d_z(p^{u+a})| \leq |d_z(p^a)||d_z(p^u)|$,

$$\begin{aligned} \sum_{a=0}^{\infty} \frac{|d_z(p^a)|^2}{p^{2a}} + 2 \sum_{u=1}^{\infty} \frac{1}{p^{u(1-\alpha)}} \sum_{a=0}^{\infty} \frac{|d_z(p^a)d_z(p^{u+a})|}{p^{2a}} &\leq \sum_{a=0}^{\infty} \frac{|d_z(p^a)|^2}{p^{2a}} \left(1 + 2 \sum_{u=1}^{\infty} \frac{|d_z(p^u)|}{p^{u(1-\alpha)}}\right) \\ (4.3) \quad &\leq \sum_{a=0}^{\infty} \frac{|d_z(p^a)|^2}{p^{2a}} \left(2 \left(1 - \frac{\delta}{p^{1-\alpha}}\right)^{-\delta k} - 1\right). \end{aligned}$$

Now consider the case $r = 0$ and note that $k \leq p$ for all $p \in I_0$. Here we use the bound (4.3) in (4.1). We choose $\alpha = 1/(10 \log_2 T)$ and note that for $p \in I_0$, $2(1 - \delta/p^{1-\alpha})^{-\delta k} - 1 \leq 2(1 - e^{1/9}/p)^{-k} - 1 \leq e^{4k/p}$. Hence we get that

$$N_0 \leq D_0 \exp \left(-\frac{\log M_0}{10 \log_2 T} + 4k \sum_{k \leq p \leq y} \frac{1}{p} \right) \leq D_0 \exp \left(-\frac{\log M_0}{10 \log_2 T} + \frac{4k}{\log k} \sum_{k \leq p \leq y} \frac{\log p}{p} \right).$$

Now $\sum_{k \leq p \leq y} \log p/p \leq \log(25y/k)$ (see Theorem I.1.7 of Tenenbaum [7]) and recall that $k \leq \log T/(e^{10} \log(y/\log T))$ and that $M_0 = T^{1/5}$. The bound in the lemma then follows in this case.

Suppose now that $r \geq 1$ so that $p \leq k$ for all $p \in I_r$. Here we use the bound (4.2) in (4.1). We take $\alpha = 1/(10 \cdot 2^{r/2} \log(ek))$ and note that for $p \leq k$,

$$\begin{aligned} \left(1 - \frac{\delta}{p^{1-\alpha}}\right)^{-\delta} \left(1 - \frac{\delta}{p^{1+\alpha}}\right)^{-\delta} \left(1 - \frac{\delta}{p}\right)^{2\delta} &\leq \left(1 - \frac{p(p^\alpha + p^{-\alpha} - 2)}{(p-1)^2}\right)^{-1} \\ &\leq \exp \left(\frac{\log^2 p}{10 \cdot 2^r p \log^2(ek)} \right). \end{aligned}$$

Using also the lower bound in Lemma 4.3 we obtain that

$$(4.4) \quad N_r \leq D_r \exp \left(-\frac{\log M_r}{10 \cdot 2^{r/2} \log(ek)} + \sum_{p \in I_r} \left(\log \frac{100k}{p} + \frac{k \log p}{10 \cdot 2^r p \log(ek)} \right) \right).$$

If $1 \leq r \leq J-1$ then we deduce that

$$\begin{aligned} N_r &\leq D_r \exp \left(-\frac{\log M_r}{10 \cdot 2^{r/2} \log(ek)} + \sum_{k/2^r \leq p \leq k/2^{r-1}} (r+5) \right) \\ &\leq D_r \exp \left(-\frac{\log M_r}{10 \cdot 2^{r/2} \log(ek)} + \frac{8(r+5)k}{2^r \log(ek)} \right) \end{aligned}$$

and since $\log M_r = (\log T)/(5r^2)$ this gives $N_r \leq D_r \exp(-\log T/(\log_2 T)^4)$ for large T . If $r = J$ and $k \leq 10^6$ then the Lemma follows at once from (4.4). If $r = J$ and $k > 10^6$ then (4.4) gives that

$$\begin{aligned} N_r &\leq D_r \exp \left(-\frac{\log M_J}{10 \cdot 2^{J/2} \log(ek)} + \sum_{p \leq k/(\log k)^4} \left(\log \frac{100k}{p} + \frac{k \log p}{10 \cdot 2^J p \log(ek)} \right) \right) \\ &\leq D_r \exp \left(-\frac{\log M_J}{10 \cdot 2^{J/2} \log(ek)} + O\left(\frac{\log T}{(\log_2 T)^4} \right) \right), \end{aligned}$$

which proves the Lemma in this case.

Proof of Proposition 4.2. We prove Proposition 4.2 by induction on the cardinality of R . The case when $R = \emptyset$ is clear and suppose the Proposition holds for all proper subsets of R . We expand

$$|\zeta(1+it; R)|^{2z} = \sum_{\substack{m_r, n_r \in \mathcal{S}(\{r\}) \\ \text{for all } r \in R}} \prod_{r \in R} \left(\frac{d_z(m_r) d_z(n_r)}{m_r n_r} \right) \left(\frac{\prod_{r \in R} m_r}{\prod_{r \in R} n_r} \right)^{it}.$$

Set $u_r = m_r n_r / (m_r, n_r)^2$. Using inclusion-exclusion we decompose the above as

$$(4.5) \quad \sum_{\substack{m_r, n_r \in \mathcal{S}(\{r\}), \text{ and} \\ u_r \leq M_r \text{ for all } r \in R}} + \sum_{\substack{W \subset R \\ W \neq \emptyset}} (-1)^{|W|-1} \sum_{\substack{m_r, n_r \in \mathcal{S}(\{r\}) \\ \text{for all } r \in R, \text{ and} \\ u_w > M_w \text{ for all } w \in W}}$$

with M_w as in Lemma 4.4.

First let us consider the contribution of the first sum in (4.5). This gives

$$(4.6) \quad \sum_{\substack{m_r, n_r \in \mathcal{S}(\{r\}), \text{ and} \\ u_r \leq M_r \text{ for all } r \in R}} \prod_{r \in R} \left(\frac{d_z(m_r) d_z(n_r)}{m_r n_r} \right) \frac{1}{T} \int_T^{2T} \left(\prod_{r \in R} \frac{m_r}{n_r} \right)^{it} dt.$$

If we reduce $\prod_{r \in R} m_r/n_r$ to lowest terms then both the numerator and denominator would be bounded by $\prod_r u_r \leq \prod_{r \in R} M_r \leq T^{\frac{(1+\pi^2/6)}{5}} \leq T^{\frac{3}{5}}$. Thus if $\prod_{r \in R} m_r/n_r \neq 1$ then

$$\frac{1}{T} \int_T^{2T} \left(\frac{\prod_{r \in R} m_r}{\prod_{r \in R} n_r} \right)^{it} dt \ll \frac{1}{T |\log \prod_r m_r/n_r|} \ll T^{-\frac{2}{5}}.$$

Hence we obtain that the expression in (4.6) equals

$$\sum_{\substack{m_r=n_r \in \mathcal{S}(\{r\}) \\ \text{for all } r \in R}} \prod_{r \in R} \left(\frac{d_z(m_r)}{m_r} \right)^2 + O\left(T^{-\frac{2}{5}} \sum_{\substack{m_r, n_r \in \mathcal{S}(\{r\}) \\ \text{for all } r \in R}} \prod_{r \in R} \left(\frac{|d_z(m_r)d_z(n_r)|}{m_r n_r} \right)\right).$$

The main term above is $\sum_{n \in \mathcal{S}(R)} d_z(n)^2/n^2$. The error term is $\ll T^{-\frac{2}{5}} \prod_{p \in \cup_{r \in R} I_r} (1 - \delta/p)^{-2k\delta}$ and using the lower bound of Lemma 4.3 this is $\ll T^{-\frac{1}{3}} \sum_{n \in \mathcal{S}(R)} d_z(n)^2/n^2$. Thus the contribution of the first term in (4.5) is

$$(4.7) \quad (1 + O(T^{-\frac{1}{3}})) \sum_{n \in \mathcal{S}(R)} \frac{d_z(n)^2}{n^2}.$$

Now we consider the contribution of the second term in (4.5). This gives

$$\begin{aligned} & \sum_{\substack{W \subset R \\ W \neq \emptyset}} (-1)^{|W|-1} \sum_{\substack{m_w, n_w \in \mathcal{S}(\{w\}), \text{ and} \\ u_w > M_w \text{ for all } w \in W}} \prod_{w \in W} \left(\frac{d_z(m_w)d_z(n_w)}{m_w n_w} \right) \\ & \times \frac{1}{T} \int_T^{2T} \left(\frac{\prod_{w \in W} m_w}{\prod_{w \in W} n_w} \right)^{it} |\zeta(1+it; R-W)|^{2z} dt, \end{aligned}$$

which is bounded in magnitude by

$$\sum_{\substack{W \subset R \\ W \neq \emptyset}} \sum_{\substack{m_w, n_w \in \mathcal{S}(\{w\}), \text{ and} \\ u_w > M_w \text{ for all } w \in W}} \prod_{w \in W} \left(\frac{|d_z(m_w)d_z(n_w)|}{m_w n_w} \right) \frac{1}{T} \int_T^{2T} |\zeta(1+it; R-W)|^{2z} dt.$$

By the induction hypothesis we see that

$$\frac{1}{T} \int_T^{2T} |\zeta(1+it; R-W)|^{2z} dt \ll \sum_{n \in \mathcal{S}(R-W)} \frac{d_z(n)^2}{n^2},$$

while from Lemma 4.4 (with $m = u_w$ and $\ell = (m_w, n_w)$ so that $d_z(m\ell)d_z(\ell) = d_z(m_w)d_z(n_w)$; and note that the number of pairs m_w, n_w which give rise to a given pair ℓ, m is exactly $2^{\omega(m)}$) we deduce that

$$\sum_{\substack{m_w, n_w \in \mathcal{S}(\{w\}) \\ u_w > M_w}} \frac{|d_z(m_w)d_z(n_w)|}{m_w n_w} \leq \sum_{n \in \mathcal{S}(\{w\})} \frac{d_z(n)^2}{n^2} \exp\left(-\frac{\log T}{(\log_2 T)^4}\right).$$

From these estimates it follows that the contribution of the second term in (4.5) is

$$\ll |R| \sum_{n \in \mathcal{S}(R)} \frac{d_z(n)^2}{n^2} \exp\left(-\frac{\log T}{(\log_2 T)^4}\right).$$

Combining this with (4.7) we obtain Proposition 4.2.

Proof of Theorem 4.1. In view of Proposition 4.2 it remains only to prove that

$$(4.8) \quad \sum_{\substack{n=1 \\ p|n \Rightarrow p \leq y}}^{\infty} \frac{d_z(n)^2}{n^2} = \prod_{p \leq k} \left(1 - \frac{\delta}{p}\right)^{-2k\delta} \exp\left(\frac{2k}{\log k} \left(C + O\left(\frac{k}{y} + \frac{1}{\log k}\right)\right)\right).$$

Using the first part of Lemma 4.3 for $p \geq \sqrt{k}$ and the second part for $p < \sqrt{k}$ we see that

$$\sum_{\substack{n=1 \\ p|n \Rightarrow p \leq y}}^{\infty} \frac{d_z(n)^2}{n^2} = \prod_{p < \sqrt{k}} \left(1 - \frac{\delta}{p}\right)^{-2k\delta} \prod_{\sqrt{k} \leq p \leq y} I_0\left(\frac{2k}{p}\right) \exp(O(\sqrt{k})).$$

Since $\log I_0(t) = O(t^2)$ for $0 \leq t \leq 2$ we have by the prime number theorem and partial summation that

$$\begin{aligned} \sum_{k \leq p \leq y} \log I_0\left(\frac{2k}{p}\right) &= \frac{2k}{\log k} \int_{2k/y}^2 \log I_0(t) \frac{dt}{t^2} + O\left(\frac{k}{\log^2 k}\right) \\ &= \frac{2k}{\log k} \int_0^2 \log I_0(t) \frac{dt}{t^2} + O\left(\frac{k^2}{y \log k} + \frac{k}{\log^2 k}\right). \end{aligned}$$

Since $\log I_0(t) = t + O(\log t)$ for $t \geq 2$ we obtain by the prime number theorem and partial summation that

$$\sum_{\sqrt{k} \leq p \leq k} \left(\log I_0\left(\frac{2k}{p}\right) + 2k\delta \log\left(1 - \frac{\delta}{p}\right)\right) = \frac{2k}{\log k} \int_2^{\infty} (\log I_0(t) - t) \frac{dt}{t^2} + O\left(\frac{k}{\log^2 k}\right).$$

These estimates prove (4.8) and so Theorem 4.1 follows.

5. PROOF OF THEOREM 1

Let $\log T(\log_2 T)^4 \geq y \geq e^2 \log T$, and let $T\Phi_T(\tau; y)$ denote the measure of points $t \in [T, 2T]$ for which $|\zeta(1+it; y)| \geq e^\gamma \tau$. Taking $z = k$ for an integer $3 \leq k \leq \log T/(e^{10} \log(y/\log T))$ in Theorem 4.1 and using Mertens' theorem $\prod_{p \leq k} (1 - 1/p)^{-1} = e^\gamma \log k + O(1/\log^2 k)$ we get that

$$\begin{aligned} 2k \int_0^{\infty} \Phi_T(t; y) t^{2k-1} dt &= \frac{1}{T} \int_T^{2T} e^{-2k\gamma} |\zeta(1+it; y)|^{2k} dt \\ (5.1) \quad &= (\log k)^{2k} \exp\left(\frac{2k}{\log k} \left(C + O\left(\frac{k}{y} + \frac{1}{\log k}\right)\right)\right). \end{aligned}$$

Now $\int_0^\infty \Phi_T(t; y) dt = e^{-\gamma}(1/T) \int_T^{2T} |\zeta(1+it; y)| dt \leq e^{-\gamma}((1/T) \int_T^{2T} |\zeta(1+it; y)|^4 dt)^{1/4} \ll 1$ by Theorem 4.1; so, by Hölder's inequality,

$$\int_0^\infty \Phi_T(t; y) t^a dt \leq \left(\int_0^\infty \Phi_T(t; y) dt \right)^{1-a/b} \left(\int_0^\infty \Phi_T(t; y) t^b dt \right)^{a/b} \ll \left(\int_0^\infty \Phi_T(t; y) t^b dt \right)^{a/b}$$

for $a < b$. While (5.1) at present holds only for integer values of k , we may interpolate to non-integer value $\kappa \in (k-1, k)$ by taking $a = 2k-3$, $b = 2\kappa-1$ and then $a = 2\kappa-1$, $b = 2k-1$ in the last inequality to obtain

$$\left(\int_0^\infty \Phi_T(t; y) t^{2k-3} dt \right)^{\frac{2\kappa-1}{2k-3}} \ll \int_0^\infty \Phi_T(t; y) t^{2\kappa-1} dt \ll \left(\int_0^\infty \Phi_T(t; y) t^{2k-1} dt \right)^{\frac{2\kappa-1}{2k-1}},$$

and so we get (5.1) for κ by substituting (5.1) for $k-1$ and k into this equation.

Suppose $1 \ll \tau \leq \log_2 T - 20 - \log_2(y/\log T)$ and select $\kappa = \kappa_\tau$ such that $\log \kappa = \tau - 1 - C$. Let $\epsilon > 0$ be a bounded parameter to be fixed shortly and put $K = \kappa e^\epsilon$. Observe that

$$\begin{aligned} 2k \int_{\tau+\epsilon}^\infty \Phi_T(t; y) t^{2\kappa-1} dt &\leq 2k(\tau + \epsilon)^{2\kappa-2K} \int_{\tau+\epsilon}^\infty \Phi_T(t; y) t^{2K-1} dt \\ &\leq (\tau + \epsilon)^{2\kappa(1-e^\epsilon)} \left(2K \int_0^\infty \Phi_T(t; y) t^{2K-1} dt \right). \end{aligned}$$

Using (5.1) we deduce that

$$\begin{aligned} &2K \int_0^\infty \Phi_T(t; y) t^{2K-1} dt \\ &= \left((\log \kappa + \epsilon) \exp \left(\frac{C}{\log \kappa} \left(1 + O \left(\frac{1}{\log \kappa} + \frac{\kappa}{y} \right) \right) \right) \right)^{2K} \\ &= \exp \left(\frac{2\kappa(\epsilon e^\epsilon + C(e^\epsilon - 1))}{\log \kappa} + O \left(\frac{\kappa}{\log^2 \kappa} + \frac{\kappa^2}{y \log \kappa} \right) \right) (\log \kappa)^{2\kappa(e^\epsilon - 1)} \int_0^\infty \Phi_T(t; y) t^{2\kappa-1} dt. \end{aligned}$$

We conclude from the above that

$$2\kappa \int_{\tau+\epsilon}^\infty \Phi_T(t; y) t^{2\kappa-1} dt = \exp \left(\frac{2\kappa}{\log \kappa} (1 + \epsilon - e^\epsilon) + O \left(\frac{\kappa}{\log^2 \kappa} + \frac{\kappa^2}{y \log \kappa} \right) \right) \int_0^\infty \Phi_T(t; y) t^{2\kappa-1} dt.$$

Choose $\epsilon = c(1/\tau + (\log T)/y)^{\frac{1}{2}}$ for a suitable constant $c > 0$, so that for large τ (and hence large κ),

$$\int_{\tau+\epsilon}^\infty \Phi_T(t; y) t^{2\kappa-1} dt \leq \frac{1}{100} \int_0^\infty \Phi_T(t; y) t^{2\kappa-1} dt,$$

say. A similar argument reveals that

$$\int_0^{\tau-\epsilon} \Phi_T(t; y) t^{2\kappa-1} dt \leq \frac{1}{100} \int_0^\infty \Phi_T(t; y) t^{2\kappa-1} dt.$$

Combining these two assertions with (5.1) for κ we obtain

$$\int_{\tau-\epsilon}^{\tau+\epsilon} \Phi_T(t; y) t^{2\kappa-1} dt = (\log \kappa)^{2\kappa} \exp\left(\frac{2\kappa C}{\log \kappa} (1 + O(\epsilon^2))\right).$$

Since Φ_T is a non-increasing function we deduce that the left side above is

$$\geq \Phi_T(\tau + \epsilon; y) \tau^{2\kappa} \exp(O(\kappa\epsilon/\tau)), \quad \text{and} \quad \leq \Phi_T(\tau - \epsilon; y) \tau^{2\kappa} \exp(O(\kappa\epsilon/\tau)).$$

It follows that

$$\Phi_T(\tau + \epsilon; y) \leq \exp\left(- (2 + O(\epsilon)) \frac{e^{\tau-1-C}}{\tau}\right) \leq \Phi_T(\tau - \epsilon; y),$$

and hence that uniformly in $\tau \leq \log_2 T - 20 - \log_2(y/\log T)$ we have

$$(5.2) \quad \Phi_T(\tau; y) = \exp\left(- \frac{2e^{\tau-1-C}}{\tau} (1 + O(\epsilon))\right).$$

From Proposition 3.2 we know that $\Phi_T(\tau) = \Phi_T(\tau + O(\epsilon); y) + O(\exp(-\log T/50 \log_2 T))$ for $\tau \ll \log_2 T$; and so from (5.2) we deduce that uniformly in $\tau \leq \log_2 T - 20 - \log(y/\log T)$ we have

$$\Phi_T(\tau) = \exp\left(- \frac{2e^{\tau-1-C}}{\tau} (1 + O(\epsilon))\right) + O\left(\exp\left(- \frac{\log T}{50 \log_2 T}\right)\right).$$

Taking $y = \min(\tau \log T, (\log^2 T)/e^{10+\tau})$ above we easily obtain Theorem 1 for Φ_T . The argument for Ψ_T is analogous, using $z = -k$ in Theorem 4.1.

One finds, using the first part of Lemma 4.3 and the observation that $\log I_0(2k/p) \ll k^2/p^2$ for $p > k$, that

$$\begin{aligned} \mathbb{E}(|L(1, X)|^{2z}) &= \sum_{n \geq 1} \frac{d_z(n)^2}{n^2} = \sum_{\substack{n=1 \\ p|n \implies p \leq y}}^{\infty} \frac{d_z(n)^2}{n^2} \exp\left(O\left(\frac{k^2}{y \log y}\right)\right) \\ &= \prod_{p \leq k} \left(1 - \frac{\delta}{p}\right)^{-2k\delta} \exp\left(\frac{2k}{\log k} \left(C + O\left(\frac{k}{y} + \frac{1}{\log k}\right)\right)\right), \end{aligned}$$

the last line following as in the proof of Theorem 4.1. With this estimate we can proceed precisely as in the proof of Theorem 1 to obtain the analogous estimate.

6. LARGE VALUES OF $|\zeta(1 + it)|$: PROOF OF THEOREM 2

Let T be large and put $y = \log T \log_2 T / (4B \log_3 T)$ for some $B \geq 5$, and $\delta = 1/[\log_2 T]^4$. Let $\|z\|$ denote the distance of z from the nearest integer.

Lemma 6.1. *For any real t_0 there is a positive integer $m \leq T^{\frac{1}{B}}$ such that for each prime $p \leq y$ we have $\| (mt_0 \log p)/2\pi \| \leq \delta$.*

Proof. This follows from Dirichlet's theorem on Diophantine approximation (see for example §8.2 of [8]) since $1/\delta$ is an integer and $(1/\delta)^{\pi(y)} \leq T^{\frac{1}{B}}$, by the prime number theorem.

Lemma 6.2. *For any real t_1 there is a positive integer $n \leq [\log_2 T]^2$ for which*

$$\operatorname{Re} \sum_{y \leq p \leq \exp((\log T)^{10})} \frac{1}{p^{1+int_1}} \geq -\frac{10}{\log_2 T}.$$

Proof. Let $K(x) = \max(0, 1 - |x|)$ and note that $\sum_{l=-L}^L K(l/L)e^{ilt}$ (the Fejer kernel) is non-negative for all positive integers L and all t . It follows therefore that

$$\sum_{j=-[\log_2 T]^2}^{[\log_2 T]^2} K\left(\frac{j}{[\log_2 T]^2}\right) \sum_{y \leq p \leq \exp((\log T)^{10})} \frac{1}{p^{1+ijt_1}} \geq 0.$$

Hence we obtain that

$$\begin{aligned} \operatorname{Re} \sum_{j=1}^{[\log_2 T]^2} K\left(\frac{j}{[\log_2 T]^2}\right) \sum_{y \leq p \leq \exp((\log T)^{10})} \frac{1}{p^{1+ijt_1}} &\geq -\frac{1}{2} \sum_{y \leq p \leq \exp((\log T)^{10})} \frac{1}{p} \\ &\geq -5 \log_2 T. \end{aligned}$$

The Lemma follows at once.

Proof of Theorem 2. For $T^{\frac{1}{10}} \leq |t| \leq T$ one has

$$\log \zeta(1+it) = - \sum_{p \leq \exp((\log T)^{10})} \log \left(1 - \frac{1}{p^{1+it}}\right) + O\left(\frac{1}{\log T}\right).$$

(One can prove this, arguing as in the proof of the prime number theorem, by noting that $(1/2i\pi) \int_{(c)} \log \zeta(1+it+w)(x^w/w)dw$ with $x = \exp((\log T)^{10})$ and $c > 0$ gives the main term of the right side by Perron's formula, and by shifting the contour to the left of 0, but enclosing a region free of zeros of $\zeta(s)$, we get residue $\log \zeta(1+it)$ from the simple pole at $w = 0$, and the error term from the remaining integral.)

Combining Lemmas 6.1 and 6.2 (with $t_1 = mt_0$) we see that for any $t_0 \in [T^{1/10}, T]$ there exists an integer ℓ (where $\ell = mn$) with $1 \leq \ell \leq T^{\frac{1}{B}} [\log_2 T]^2$ such that $\| (\ell t_0 \log p)/2\pi \| \leq 1/[\log_2 T]^2$ for each prime $p \leq y$, and such that

$$\operatorname{Re} \sum_{y \leq p \leq \exp((\log T)^{10})} \frac{1}{p^{1+i\ell t_0}} \geq -\frac{10}{\log_2 T}.$$

We deduce therefore that

$$\begin{aligned} |\zeta(1 + it_0)| &\geq \prod_{p \leq y} \left(1 - \frac{1}{p} + O\left(\frac{1}{p(\log_2 T)^2}\right)\right)^{-1} \left(1 + O\left(\frac{1}{\log_2 T}\right)\right) \\ &\geq e^\gamma (\log_2 T + \log_3 T - \log_4 T - \log A + O(1)), \end{aligned}$$

using the prime number theorem, where $A = 1/(2/B + 3 \log_2 T / \log T)$.

We use the above procedure with $t_0 = T_0, T_0 + 1, T_0 + 2, \dots, T_0 + U_0$ where $T_0 = [T^{1-1/B}/3[\log_2 T]^2]$ and $U_0 = [T^{1-2/B}/7[\log_2 T]^4]$. Let ℓ_i be as above so $\ell_i \leq T^{1/B}[\log_2 T]^2$ and thus $\tau_i = \ell_i(T_0 + i) \leq T/2$. We claim that $|\tau_i - \tau_j| \geq 1$ if $i \neq j$ for if not then evidently $\ell_i \neq \ell_j$ (else $1 \leq |(T_0 + j) - (T_0 + i)| = |\tau_j - \tau_i|/\ell_i < 1$), so that

$$T_0 \leq |(\ell_i - \ell_j)T_0| \leq |\tau_i - \tau_j| + |i\ell_i - j\ell_j| < 1 + U_0 T^{1/B}[\log_2 T]^2,$$

which is false. Now each $|\zeta(1 + i\tau_j)| \geq e^\gamma (\log_2 T + \log_3 T - \log_4 T - \log A + O(1))$. Since $|\zeta'(1 + it)| \ll \log^2 T$ for $1 \leq |t| \leq T$ we see that for any $|\alpha| \leq 1/\log^2 T$ we have that $|\zeta(1 + i\tau_j + i\alpha)| = |\zeta(1 + i\tau_j)| + O(\alpha \log^2 T) = |\zeta(1 + i\tau_j)| + O(1)$. Thus the measure of $t \in [0, T]$ with $|\zeta(1 + it)| \geq e^\gamma (\log_2 T + \log_3 T - \log_4 T - \log A + O(1))$ is at least $2U_0/\log^2 T$, proving Theorem 2.

7. THE ANALOGOUS RESULTS FOR L -FUNCTIONS AT 1

By analogous methods one can prove:

Theorem 3. *Let q be a large prime.*

(i) *The proportion of characters $\chi \pmod{q}$ for which $|L(1, \chi)| > e^\gamma \tau$ is*

$$(7.1) \quad \exp\left(-\frac{2e^{\tau-C-1}}{\tau}\left(1 + O\left(\frac{1}{\tau^{\frac{1}{2}}} + \left(\frac{e^\tau}{\log q}\right)^{\frac{1}{2}}\right)\right)\right),$$

uniformly in the range $1 \ll \tau \leq \log_2 q - 20$. The same asymptotic also holds for the proportion of characters $\chi \pmod{q}$ for which $|L(1, \chi)| < \pi^2/6e^\gamma \tau$.

(ii) *There are at least $q^{1-1/A}$ characters $\chi \pmod{q}$ such that*

$$|L(1, \chi)| \geq e^\gamma (\log_2 q + \log_3 q - \log_4 q - \log A + O(1)),$$

for any $A \geq 10$.

If, in addition, we vary over all characters $\chi \pmod{q}$ and all primes $Q \leq q \leq 2Q$, then we can get a good estimate for the distribution function of $|L(1, \chi)|$ in almost the entire viable range. Thus we may prove that the proportion of $|L(1, \chi)| \geq e^\gamma \tau$ is (7.1) for the range $1 \leq \tau \leq \log_2 Q + \log_3 Q - 100$, but now with the error term “ $(e^\tau/(\log Q \log_2 Q))^{\frac{1}{2}}$ ” in place of “ $(e^\tau/\log q)^{\frac{1}{2}}$ ” (and a corresponding result holds for $1/|(6/\pi^2)L(1, \chi)|$).

The broad outline of the proof is the same, though now replacing $\log T$ by $\log Q \log_2 Q$, so that $\log Q(\log_2 Q)^4 \geq y \geq e^2 \log Q \log_2 Q$ and the range for k becomes $2 \leq k \leq$

$\log Q \log_2 Q / (e^{10} \log(y/(\log Q \log_2 Q)))$. The result follows easily from the following analogy to Theorem 4.1,

$$\frac{1}{\pi(Q)} \sum_{q \leq Q} \frac{1}{\varphi(q)} \sum_{\chi \pmod{q}} |L(1, \chi; y)|^{2z} = \prod_{p \leq k} \left(1 - \frac{\delta}{p}\right)^{-2k\delta} \exp\left(\frac{2k}{\log k} \left(C_1 + O\left(\frac{k}{y} + \frac{1}{\log k}\right)\right)\right),$$

and an appropriate development of Lemma 4.3, where $L(1, \chi; y) := \prod_{p \leq y} (1 - \chi(p)/p)^{-1}$. The above estimate, though, is proved rather more easily than Theorem 4.1. Since $L(1, \chi; y)^z = \sum_{n \in \mathcal{S}(y)} d_z(n) \chi(n)/n$, and $L(1, \bar{\chi}; y)^z = \sum_{m \in \mathcal{S}(y)} d_z(m) \bar{\chi}(m)/m$ where $\mathcal{S}(y)$ is the set of integers all of whose prime factors are $\leq y$, the left side of this equation equals

$$\sum_{m, n \in \mathcal{S}(y)} \frac{d_z(m) d_z(n)}{mn} \left\{ \frac{1}{\pi(Q)} \sum_{q \leq Q} \frac{1}{\varphi(q)} \sum_{\chi \pmod{q}} \chi(m) \bar{\chi}(n) \right\}.$$

The term in $\{\}$ equals $1 - \#\{q \leq Q : q|mn\}/\pi(Q)$ if $m = n$, and is $\leq \#\{q \leq Q : q|m - n\}/\pi(Q)$ if $m \neq n$. Therefore our sum is

$$\sum_{n \in \mathcal{S}(y)} \frac{d_z(n)^2}{n^2} + O\left(\frac{1}{\pi(Q)} \left(\sum_{m \in \mathcal{S}(y)} \frac{|d_z(m)| \log 2m}{m} \right)^2\right).$$

Now $\log 2n \ll k^2 + n^{1/k}$ so that

$$\begin{aligned} \sum_{n \in \mathcal{S}(y)} \frac{|d_z(n)|}{n} \log 2n &\ll k^2 \prod_{p \leq y} \left(1 - \frac{\delta}{p}\right)^{-\delta k} + \prod_{p \leq y} \left(1 - \frac{\delta}{p^{1-1/k}}\right)^{-\delta k} \\ &\ll \prod_{p \leq y} \left(1 - \frac{\delta}{p}\right)^{-\delta k} \left(k^2 + \exp\left(O\left(k \sum_{p \leq y} \frac{p^{1/k} - 1}{p}\right)\right)\right) \\ &\ll (\log Q)^{O(1)} \prod_{p \leq y} \left(1 - \frac{\delta}{p}\right)^{-\delta k}, \end{aligned}$$

and the claimed estimate follows from Lemma 4.3.

REFERENCES

1. R. Balasubramanian, K. Ramachandra, and A. Sankaranarayanan, *On the frequency of Titchmarsh's phenomenon for $\zeta(s)$ -VIII*, Proc. Ind. Acad. Sci. **102** (1992), 1-12.
2. A. Granville and K. Soundararajan, *The distribution of values of $L(1, \chi_d)$* , Geometric and Funct. Anal. **13** (2003), 992-1028.
3. N. Levinson, *Ω -theorems for the Riemann zeta-function*, Acta Arith. **20** (1972), 319-332.
4. J.E. Littlewood, *On the function $1/\zeta(1+it)$* , Proc. London Math. Soc. **27** (1928), 349-357.
5. J.E. Littlewood, *On the class number of the corpus $P(\sqrt{-k})$* , Proc. London Math. Soc. **27** (1928), 358-372.
6. K. Ramachandra, *On the frequency of Titchmarsh's phenomenon for $\zeta(s)$ - VII*, Ann. Acad. Sci. Fenn. **14** (1989), 27-40.
7. G. Tenenbaum, *Introduction to analytic and probabilistic number theory*, Cambridge Studies in Advanced Mathematics **46**, Cambridge University Press, Cambridge, 1995.

8. E.C. Titchmarsh, *The theory of the Riemann zeta-function*, Oxford University Press, Oxford, 1986.
9. I.M. Vinogradov, *A new estimate for $\zeta(1+it)$* , Izv. Akad. Nauk SSSR Ser. Mat **22** (1958), 161–164.

DÉPARTEMENT DE MATHÉMATIQUES ET STATISTIQUE, UNIVERSITÉ DE MONTRÉAL, CP 6128 SUCC
CENTRE-VILLE, MONTRÉAL, QC H3C 3J7, CANADA

E-mail address: `andrew@dms.umontreal.ca`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MICHIGAN 48109, USA

E-mail address: `ksound@umich.edu`